European Journal of Innovation in Nonformal Education (EJINE)
Volume 6 | Issue 2 | Feb - 2026 ISSN: 2795-8612

Interrelationships Between Anaerobic Power, Muscle Fiber Type Ratio, and Start Reaction Time in Short-Distance Sprinting Events (100 M and 200 M) in Track and Field Athletics

Egamberdiev Ilhom Ergashevich NamDU
Professor of the Department of Sports Activity and Physical Culture

A B S T R A C T

The connection between anaerobic power, muscle fiber makeup, and reaction time at the start is crucial in influencing performance results in short-distance sprints like the 100 m and 200 m in athletics. This article examines the physiological bases of these elements, highlighting how anaerobic energy systems, mainly phosphagen and glycolytic pathways, interact with the dominance of fast-twitch muscle fibers to improve explosive power production and swift neural activation essential for peak starting performance. Utilizing biomechanical and bioenergetic viewpoints, it explores the hereditary and adaptable components of muscle fiber allocation, the energy requirements of sprinting, and the neuromuscular processes that control reaction time. The analysis indicates significant correlations showing that athletes with a greater percentage of Type II muscle fibers demonstrate enhanced anaerobic power and faster reaction times, aiding in improved acceleration and speed retention. This synthesis highlights the interconnectedness of these components in elite sprinting, providing perspectives on training implications for improving sprint-specific adaptations while maintaining overall athletic growth.

A R T I C L E I N F O

Article history:

Received 16 Dec 2025
Received in revised form 16 Jan 2026
Accepted 08 Feb 2026

Keywords: Anaerobic power, muscle fiber composition, start reaction time, sprint performance, track and field, 100 m sprint, 200 m sprint, fast-twitch fibers, phosphagen system, glycolytic pathway, neuromuscular activation.

Hosting by Innovatus Publishing Co. All rights reserved. © 2026

In track and field athletics, especially in the 100 m and 200 m sprints, the relationship between anaerobic power, muscle fiber type, and reaction time at the start creates a crucial trio that influences an athlete's capacity for peak performance. Anaerobic power denotes the body's ability to produce energy quickly without using oxygen, mainly through the phosphagen and glycolytic systems. This is crucial in sprints where the time frame is too short for considerable aerobic input, usually lasting less than 10-20 seconds. For example, in a 100 m sprint, the relative anaerobic contribution can be as high as 79-91% based on oxygen deficit measurements, with the phosphagen system supplying immediate high-energy phosphates for the initial surge and glycolysis supporting the effort via lactate production. Athletes possessing increased anaerobic power are capable of generating greater force outputs in less time, facilitating rapid acceleration from the starting blocks. The composition of muscle fibers greatly influences this anaerobic capability. Human skeletal muscles consist of a combination of slow-twitch (Type I) and fast-twitch (Type IIa and IIx) fibers, with the fast-twitch fibers being more prevalent in top sprinters. Type II fibers are distinguished by their elevated myosin ATPase activity, increased cross-sectional areas, and larger reserves of phosphocreatine, enabling swift contraction and force production.

Research indicates that sprinters typically have 50-70% Type II fibers in important muscles such as the vastus lateralis, while endurance athletes might carry just 30-40%. This structure not only boosts maximum power output to levels of 30-36 W/kg in elite sprinters but also relates to the capacity to sustain speed in the face of fatigue. The proportion of cross-sectional area taken up by Type II fibers serves as a more reliable predictor of sprinting ability than just fiber percentage or total area, as it indicates the functional capability for intense effortsA larger proportion of Type IIx fibers is associated with enhanced jumping and strength performance, which directly impacts sprint acceleration phases.

The onset of reaction time, which is the period from the signal given by the starter to the initial movement, is another neuromuscular factor closely connected to these components. Usually between 0.17-0.18 seconds in top athletes, it includes sensory processing, neural transmission, and recruitment of motor units. Quicker reaction times relate to improved neuromuscular efficiency, in which the swift activation of alpha motor neurons prompts instant muscle contraction. In this context, the dominance of fast-twitch fibers serves a dual purpose: they facilitate rapid force generation while also aligning with genetic tendencies for short-latency neural pathways. Correlations suggest that athletes who have higher Type II fiber ratios demonstrate faster reaction times, possibly owing to increased conduction speeds in larger motor axon diameters and improved excitation-contraction coupling from calcium release in the sarcoplasmic reticulum.

The connections between these elements are complex and mutually reinforcing. Anaerobic power is enhanced by a muscle fiber composition that favors Type II subtypes, since these fibers are proficient in anaerobic ATP resynthesis, enabling prolonged high-power performance during the acceleration stage of a sprint. In the 100 m, where peak speed is attained between 40-60 m, this cooperation allows athletes to traverse the distance using fewer strides at increased frequencies. Likewise, in the 200 m, which requires a bit more stamina because of the curve and longer time, the glycolytic aspect is more significant, but it continues to depend on the basic anaerobic strength enhanced by fast-twitch proficiency. Reaction time factors into this dynamic by establishing the initial momentum; a lag of just 0.01 seconds can result in a 0.1 m disadvantage at peak speeds surpassing 12 m/s. Neuromuscular training adaptations, including plyometric and resistance training, can strengthen these connections by boosting Type II fiber activation and hypertrophic growth, thus enhancing the rate of force production, a crucial measure where sprinters excel over endurance athletes by 20-30%. From a bioenergetic perspective, the phosphagen system takes precedence during the initial 1-2 seconds of a sprint, coinciding seamlessly with the reaction phase at the start. In this timeframe, creatine kinase quickly synthesizes ATP from phosphocreatine, powered by the abundant phosphagen reserves in Type II fibers. As the sprint continues, glycolysis dominates, creating ATP through substrate-level phosphorylation and forming lactate. Athletes possessing enhanced muscle fiber compositions can manage elevated lactate levels reaching 12-15 mmol/L after sprints without experiencing severe acidosis, due to improved buffering abilities and enzyme functions such as lactate dehydrogenase in fast-twitch fibers. This metabolic resilience directly impacts the recovery of reaction times in repeated efforts, as demonstrated in training protocols that include high-intensity interval training, simulating race conditions.

Genetic elements form the basis of this connection, with variations in genes such as ACTN3 (which encodes alpha-actinin-3, found commonly in fast-twitch fibers) linked to sprinting ability. Although the distribution of fiber types is primarily determined by genetics and is only slightly influenced by training, hypertrophic workouts can boost the cross-sectional area of Type II fibers by 10-20%, improving power production. Environmental factors like early focus on sprint training enhance these characteristics by boosting neural drive and synchronization of motor units, which subsequently reduce reaction times via practice-induced plasticity in the central nervous system.

Training implications emphasize the necessity for cohesive programs that address all three components. For anaerobic power, brief sprints (20-40 m) with complete recovery focus on restoring phosphagen levels, whereas longer intervals (60-100 m) enhance glycolytic tolerance. To enhance muscle fiber usage, strength training with heavy weights (80-90% of 1RM) encourages Type II growth and strength. Drills involving reaction time, featuring auditory and visual cues in initial positions, boost neuromuscular responsiveness, with data indicating enhancements of 0.02-0.05 seconds with regular practice.

High-intensity warm-up protocols can enhance anaerobic performance by 8-10%, effectively preparing the phosphagen system for explosive movements. In elite environments, these connections appear as differences in performance. Elite 100 m sprinters, recording times under 9.8 seconds, commonly exhibit anaerobic power outputs surpassing 30 W/kg, muscle fiber distributions with more than 60% Type II, and reaction times under 0.15 seconds. In comparison, sub-elite athletes might fall short in certain areas, like inadequate Type II dominance resulting in premature fatigue or sluggish reactions exacerbating acceleration weaknesses. Longitudinal assessment through biopsy examinations, Wingate tests for anaerobic strength, and photoelectric timing for reactions can direct tailored interventions.

In the end, the unified combination of anaerobic power, muscle fiber types, and reaction time at the start highlights the overall requirements of sprinting. Improving one element consistently advantages the others; for example, boosting the proportion of Type II fibers via genetic selection or training enhances both power and reaction time. This triad elucidates differences in performances for 100 m and 200 m events while also guiding strategies for recognizing talent and optimizing conditioning, enabling athletes to reach their physiological limits in the competitive realm of track and field.

Conclusion

The complex relationship among anaerobic power, muscle fiber types, and reaction time at the start forms the essential physiological basis for achieving success in short-distance sprint events (100 m and 200 m) in track and field sports. Top performance in these fields is not dictated by any one element alone, but instead by the collaborative enhancement of all three factors. Athletes with a greater proportion of Type II (particularly Type IIx) muscle fibers show significantly enhanced anaerobic power traits, both alactic (phosphagen) and lactic (glycolytic), enabling them to produce very high horizontal forces during the acceleration phase and sustain high maximal velocity for extended periods. The same prevalence of fast-twitch fibers is strongly linked to faster neuromuscular activation sequences, leading to reduced reaction times at the onset (often under 0.150–0.160 s in elite athletes). Quicker reaction times provide a significant edge in the initial 10–20 meters, as even slight variations in milliseconds can lead to major positional advantages that are hard to regain later in the race. The connections are mutually reinforcing: an increased cross-sectional area and functional ability of Type II fibers boost both immediate power output and the rate of force production, which in turn reduces electromechanical delay and enhances reaction time.

Genetic predisposition significantly influences the maximum bounds of muscle fiber type distribution and fundamental neuromuscular efficiency, yet focused training can substantially enhance all three factors within a person's genetic range. Intense resistance workouts, plyometric exercises, sprint-specific resistance and assistance drills, along with reaction-time-centered start practice, together enhance Type II fiber growth and power generation, boost anaerobic capacity and strength, and lower both mean and variability in reaction times. In practical terms, the most accomplished sprinters of the contemporary period continually exhibit this profile:

- >60–70% type ii fiber composition in essential lower-limb muscles,
- Maximum anaerobic power levels surpassing 30–36 $w \cdot kg^{-1}$,
- Reaction times consistently under 0.160 s (typically 0.120–0.145 s),
- Along with the smooth incorporation of these attributes into a technically polished block removal and speed enhancement technique.

Consequently, short-sprint performance ought to be understood not as the addition of separate skills, but as the result of an interconnected physiological system where anaerobic energy production, muscle contraction characteristics, and neuromuscular speed enhance each other. Future progress in talent detection, personalized training strategies, and performance evaluation will increasingly depend on the concurrent assessment and enhancement of this trio rather than isolating and targeting any single element. In the end, it is the balanced enhancement of anaerobic strength, fast-twitch muscle fiber superiority, and immediate start response that distinguishes elite sprinters from other athletes.

References

1. The contribution of energy systems during 15-second sprint exercise in athletes of different sports specializations. NIH.
2. Sprinters' and Marathon Runners' Performances Are Better Explained by Muscle Fibers' Percentage Cross-Sectional Area than Any Other Parameter of Muscle Fiber Composition. MDPI.
3. Physiological and Neuromuscular Indices Associated with Sprint Running Performance. ResearchGate.
4. Relationships between muscle fibre characteristics and physical performance capacity in trained athletic boys. PubMed.
5. The Training and Development of Elite Sprint Performance: an Integration of Scientific and Best Practice Literature. Springer Nature.